

# Biorefinery of Wastes of Meat Processing for Agricultural Applications

Katalin Perei<sup>1,2</sup>, Viktória Markó<sup>1</sup>, Csilla Juhász<sup>1,2</sup>, Anett Széll<sup>1,2</sup>, Attila Bodor<sup>1,2,3</sup>, Gábor Rákhely<sup>1,2,3</sup>

<sup>1</sup>Department of Biotechnology, University of Szeged

Közép fasor 52., H-6726 Szeged, Hungary

<sup>2</sup>Institute of Environmental and Technological Sciences, University of Szeged

Közép fasor 52., H-6726 Szeged, Hungary

<sup>3</sup>Institute of Biophysics, Biological Research Centre

Temesvári krt. 62., H-6726 Szeged, Hungary

## Extended Abstract

Tremendous amount of keratin and bone wastes are generated daily in the poultry and meat industries in the world, which might be recycled as biofertilizers in agriculture.

Recycling the biogenic elements (N, P, S, Ca) present in bone, feather or hair wastes facilitates these elements to return into the biogeochemical cycle. Mineralization of bone and feather wastes might provide essential elements for the agricultural plants [1,2,3]. For several popular crops, the sulfur supplementation has also serious economical impact [4].

Two bacteria were used, *Pseudomonas* sp. F1 can mobilize the P from bone. The supernatant of this culture supplemented with keratin was augmented with *Bacillus licheniformis* KK1. After one day and a week of incubation, the protein, orto-phosphate, TOC/TN and sulfate contents of the supernatants were measured.

The supernatant of *Pseudomonas* sp. F1 contained orto-phosphate (released from bone) more than enough for the growth of *Bacillus licheniformis* KK1. The liquid containing soluble form of nitrogen, phosphorous, calcium and sulfur is an “easy to use” bio-based fertilizer targeting various cultivated plants.

Biorefinery of food wastes for production of bio-products such as biofertilizer is one of the most promising environmentally sound processes, which is not only advantageous as waste management, but also leads to reduced dependence on non-renewable sources such as phosphate rocks (in production of P fertilisers).

## Acknowledgements

The project was supported by the European Union and Hungarian State (grant agreement no. EFOP-3.6.2-16-2017-00010).

## References

- [1] L. Fernández, B. Agaras, P. Zalba, L. G. Wall and C. Valverde, “*Pseudomonas* spp. isolates with high phosphate-mobilizing potential and root colonization properties from agricultural bulk soils under no-till management,” *Biology and Fertility of Soils*, vol. 48, no. 7, pp 763–773, 2012.
- [2] K. Gurdeep and M. S. Reddy, “Effects of Phosphate-Solubilizing Bacteria, Rock Phosphate and Chemical Fertilizers on Maize-Wheat Cropping Cycle and Economics,” *Pedosphere*, vol. 25, no. 3, pp. 428–437, 2015.
- [3] L. Lange, Y. Huang, and P. K. Busk, “Microbial decomposition of keratin in nature—a new hypothesis of industrial relevance,” *Appl Microbiol Biotechnol*, vol. 100, pp. 2083–2096, 2016.
- [4] R. E. Karamanos, T. B. Goh, and D. N. Flaten, “Nitrogen and sulphur fertilizer management for growing canola on sulphur sufficient soils,” *Canadian Journal of Plant Science*, vol. 87, no. 2, pp. 201-210, 2007.