

Enhanced Phytoremediation of Petroleum Hydrocarbon-Contaminated Soil by *Chromolaena Odorata* Containing Bacteria-Endophyte

Anyasi, Raymond Oriebe^{1*}; Atagana, Harrison Ifeanyichukwu¹; Sutherland, Rene²

Department of Environmental Science, Institute for Nanotechnology and water Sustainability

¹University of South Africa

²Agricultural Research Council

eanyasro@unisa.ac.za; atagahi@unisa.ac.za; sutherlandr@arc.agric.za

+27 123376194

Abstract - The study entails an isolation of Bacterial endophyte from plant growing around a crude oil sludge dam. Based on morphological characterisation, gram reactions and 16S rDNA sequence analysis, the isolate was identified as *Bacillus sefensis* Strain CS4. Following the tests on the abiotic effects as well as the initial concentration of perylene on growth and degradation efficiency of the strain, a total degradation percentage of 89 % was observed from the initial concentration of 30 mg/l for one week. Analysis of the degradation products of perylene using GC-MS/MS, indicated a shift from the previously degradation product of other bacterial endophytes showing that the strain provided a new pathway for PAH degradation. In order to evaluate the influence of a polycyclic hydrocarbon transforming ability of the bacterial strain on the phytoremediation of petroleum aromatic hydrocarbon (PAH), *B. sefensis* was inoculated into *Chromolaena odorata* plants. The plants were grown for 16 weeks with or without PAH (500 mg/kg soil in each 1L pot) in non-sterile peat medium. Plants inoculated with the strain Ros-1 were much tolerant towards the phytotoxic effects of PAH, in terms of biomass index, leaves and stem dry weight. Although the presence of plants acted as the main effective treatment responsible for PAH dissipation (55–71%), the inoculum with the strain leads to the highest PAH removal (up to 82%). Uninoculated plant control in the contaminated soil was susceptible to the phytotoxicity of the contaminants in the parameters tested. The study therefore presented the strain as a suitable plant endophyte for enhanced phytotreatment of PAH.

Keyword: Phytoremediation, Bacterial endophytes, Petroleum aromatic hydrocarbons, *Bacillus sefensis*, *Chromolaena odorata*.

1. Introduction

Bacillus sefensis strain CS4 is an endophyte that occurs in plants growing in areas with petroleum hydrocarbon contamination. This explains the implication of the bacteria in the phytoremediation of petroleum aromatic hydrocarbon (PAH) [1]. *Pseudomonas parafulva* as an endophyte lives in a plant without causing any negative symptoms in the plant [2]. It is a Gram-negative Gammaproteobacteria belonging to the order Pseudomonadales [3-4]. The bacteria according to literature, was classified in group II, class I (Cluster 1) of the *Pseudomonas fulva* strain, on the bases of genetic recharacterization. *Bacillus species* and other bacterial endophytes have been tested in their ability for phytoremediation of organic contaminants, and in those studies have demonstrated their effectiveness in reducing or completely removing such contaminant from either soil or water [5-9].

Currently, no study has reported on the use of *B sefensis* CF4; a strain of *Bacillus* as an endophyte, let alone as an endophyte in phytoremediation studies. However, owing to the fact that there are several endophytes yet to be identified, it is imperative to test for new endophytes that could be employed in the various applications such as in fuel, medicine, environment, and agriculture. Bacteria from the genus *Bacillus* are microorganisms that effectively decompose organic pollutants through cometabolism in natural water and soil environment, hence have been used in phytoremediation applications [10]. *B sefensis* from the collection of culture was chosen for the study based on its reoccurrence in a plant-endophyte profiling study done on plants growing in the PAH-contaminated environment [11]. The high incidence of the endophyte strain generated the need for its pilot testing to establish the effectiveness of the bacteria in plants colonization and consequently in phytoremediation of PAHs.

This study used selected plants in South Africa (Siam weed). This plant has been implicated in various studies for the remediation of various soil contaminants [12-14]. Author's previous study reported the ability of *Chromolaena odorata* in

the remediation of PCB-contaminated soil [15]. Meanwhile, various other studies have reported the use of such plant in the removal of metal as well as PAHs from soil [12,16-22]. In the study by Atagana [12], *Chromolaena odorata* plant was able to extract PAHs through the root to the stem and the leaf after 90 days of exposure to a used engine oil contaminated soil. Meanwhile, a study involving endophyte enhanced phytoremediation requires that the bacteria be inoculated into the plant before the degradation reaction [23-26]. Therefore, the aim of the study was to inoculate *B. sefensis* CS4 in *Chromolaena* plant and use the inoculated plant in the remediation of PAH-contaminated soil.

2. Methodology

B. sefensis CF4 was isolated from Rye grass (*Lolium*) collected from around a crude oil sludge dam in South Africa. The interest for the strain was based on its high incidence amongst the plants sampled. A clean Rye grass was surface sterilized using 75 % (v/v) ethanol for 2 minutes, cleaned with distilled water for 1 minute and flooded with commercial bleach for 1 minute. The sterilized plant was finally washed three times using distilled water to remove the residues of the chemicals. Confirmation of the success of the sterilization was done by inoculating the water from the final rinse on an LB agar medium. The sterilized plants were separated into roots, stem and leaf and were ground using sterile mortar. The paste of the plant was streaked in bacteriological agar for three days. Single colonies were transferred into the nutrient agar and preserved. To verify the purity of the strains, a single colony was viewed under a high powered microscope [10].

Identification of the endophyte strain was done using both molecular and morphological data. Extraction of DNA was done using a commercial DNA extraction kit (Genelute DNA kit from Sigma-Aldrich). In molecular identification, PCR was used to amplify the internal transcribed spacer region of the ITS rDNA [19]. The PCR, as well as the *fragment purification* and sequencing, were performed according to Jain *et al.* (2012). Fragment similarities were compared with that of the previously published data and examined with BLASTn in GenBank. *Bacillus sefensis* CS4 was obtained from cultures maintained on potato dextrose agar (PDA: Sigma Aldrich South Africa) for 7 days at 28 °C in the dark. The bacteria were harvested and placed in test tubes containing 0.05 % (v/v) aqueous solution of Tween 20 (Merck). Suspensions were adjusted to 1×10^8 mL⁻¹ of cells of *B. sefensis* according to Weyens *et al.* [6], using a Neubauer hemocytometer.

2.1. Artificial contamination of soil

50 kg of the sterile peat was contaminated with perylene dissolved in hexane to the tune of 500 mg/kg and was *homogenised* with a manual roto. The contaminated soil was analysed to ascertain the level of contamination as the initial concentration.

2.2. Inoculation of the plant

Cuttings of *Chromolaena* were planted in 12 x 12 cm plastic pots containing potting soil at 1 cm depth. The *cuttings* were allowed to grow for three weeks. The 3 weeks old plants were each sprayed about 2 ml of the cell suspension filled with surface sterilized broth, using a sterilized plastic hand sprayer of 50 ml volume. The control experiment was equally sprayed with an equal volume of the cell-free surfactant. The entire treated and control plants were allowed for 24 hours at 25 °C, and a photoperiod of 12 hours before transplanting into another 1L pots containing 1 kg of non-sterile peat previously treated with 500 kg of the selected PAHs.

Watering of the plants in the set up was done using the manual watering system, making sure that the appreciable water is allowed into the pots. Each experiment and the control was replicated into 3 and done on three *different* dates. The experiment was allowed for the number of days depending on the allotted period. At the end of each growth period, the leaves of the plants in the experimental and control set up were harvested and were dried in the *air* on a sterile laminar flow, making sure that dead tissues were not included. About three leaves from each set up were used with 1 cm piece of each leaf being cultured in Petri dishes containing PDA with 0.1 % stick antibiotics consisting of 0.02 g each of penicillin, streptomycin, and tetracycline. The presence or absence of *B. sefensis* growth was recorded after 10 days at 25 °C.

A total of 30 plants and 90 pieces of the plant were examined, and the data expressed as colonization frequencies with *the* formula below.

2.3. Growth index

Plant growth rate was measured by means of a length of the stem (L) measured on the days of testing as L₀, L₁₀, L₂₀, and L₃₀ respectively. A control experiment was measured from the uninoculated setups. Growth indexes were then measured as (L₁-L₀)/L₀ for the length of stem and percentage of germination (presence of stem) in inoculated plants compared to the control.

The data generated were analyzed using ANOVA in excel.

3. Results

Bacillus safensis CS4 was not recorded in the entire control experiment. But the inoculation techniques were successful in establishing the bacteria as an endophyte in the sample plants, although there was a difference in the colonization frequencies based on the techniques used over time. Meanwhile, the inoculation method significantly affected the colonization of leaves in the two plants species amongst the recorded days (Table 1).




Fig. 1: Growth parameters of *C. odorata* (A/C) measured after 16-wks growing in the sterile peat substrate in the presence of perylene with *B. safensis* CS4/Tween 20 and of lengths of plant, number of mature leaves per plant and root numbers: A=Length of plant; B=Number of mature leaves; C=Root numbers.




Figure 2 (A and B): PAH concentration and percentage removal measured by GC–MS in (control): surfactant-inoculated (T20) and planted *B. safensis* CS4-inoculated (CS4) sand–peat substrate at the end of the 18-wk pot experiment. Error bars show the standard deviation. Mean values marked by the same letter are not significantly different at $P < 0.05$.

Growth parameters of *C. odorata* and tobacco plants measured after 18 wk of growth with PAHs in the growth substrate as well as in the absence (control) and in the presence of the bacterial inoculum (*Bacillus safensis* CS4) are reported in Fig. 1. As can be seen, PAH amendment exerted negative effects on uninoculated plants in terms of both biomass production and stem length. In fact, a decrease of about 65% ($P < 0.01$), 54% ($P < 0.05$) and 60% ($P < 0.01$) was measured in the biomass growth index, stem length index and root dry weight respectively, in PAH treated plants comparing with the untreated ones. On the other hand, although the infection with *B. safensis* CS4 caused a strong decrease of the biomass growth index ($<70\%$; $P < 0.01$) and above-ground dry weight ($<25\%$; $P < 0.05$) in plants grown without PAHs respect to the non-inoculated ones, DBT1 infected poplars showed root dry weight value higher than that obtained for uninoculated plants ($>32\%$, $P < 0.01$), once grown in the presence of PAHs. Nevertheless, both the biomass growth index and the stem length index seemed not to be greatly affected by *B. safensis* CS4 inoculum in tobacco grown on contaminated substrate when compared with non-inoculated plants.

Fig. 2 shows the concentrations of pyrene, chrysene and perylene at the end of the pot experiment. Regarding chrysene, it was present in very low concentrations, ranging from 0.01 to 0.26 mg kg⁻¹ substrate in the different pot sets. Due to the quite limited concentrations of PAHs in the substrate, no remarkable variation is observed in control abatement among the different trials. In planted soil–peat substrate, 3.1 ± 0.9, 4.9 ± 1.6, and 6.9 ± 1.4 mg kg⁻¹ of pyrene, chrysene and perylene, respectively, were measured. Therefore, the presence of *C. odorata* plants increased the removal of these compounds, ranging from 82% to 87% (P < 0.01), if compared to control pots. Furthermore, the infection with *P. falciparum* Ros-1 exerted a significant effect in PAH removal from the substrate. The final amount of PAHs detected in inoculated-planted soil–peat substrate was 0.27 ± 0.02 of pyrene, 0.23 ± 0.05 of chrysene and 0.34 ± 0.08 mg kg⁻¹ of perylene. Thus, the association between *C. odorata* plants and *P. falciparum* Ros-1 removed up to 99% of the total amount of PAHs amended (P < 0.05).

4. Discussion

The relationships between plants and their associated microorganisms are different and complex. Rhizospheric bacteria refer to bacteria living on the surface or around the roots, in contrast to epiphytic bacteria living on the leaves [11,24]. On the other hand, endophytic bacteria can be defined as bacteria colonizing the internal tissues of plants without causing negative effects on their host [18,25]. In the last decade, endophytic microorganisms have been taken into account for applications in assisted phytoremediation protocols. In fact, several studies have focused on the use of constructed plant-associated bacteria. On the other hand, catabolic genes and plant growth promoting mechanisms occurring in natural endophytic bacteria are also used for bioremediation purposes [26].

This study demonstrates that a bacterial isolate, originally identified for its PAH degrading capacity; *Bacillus safensis* CS4, can infest *C. odorata* plant and consequently improve the phytoremediation efficiency of PAHs. Germaine *et al.* [8] already reported the infection of nonendophytic, naphthalene-degrader strain *Pseudomonas putida* G7 in pea plant. However, the authors reported a weak phytoprotection effect of this inoculum towards plants exposed to naphthalene. This slight effect was due to the low root colonization of G7 strain (about 101 CFU g⁻¹). Conversely, in the present study *B. safensis* CS4 was shown to be capable of reaching a large population size both in PAH treated and untreated *C. odorata* roots, higher than that observed in the case of *P. putida* G7 [7-8,27].

5. Conclusion

Bacillus safensis CS4 according to this present study is found to ensure rapid and effective degradation of perylene within the plant in this case *C. odorata* despite the unavailability of existing literatures using the plant. This means that the use of endophytes of this clad to enhance the phytoremediation of petroleum hydrocarbon is encouraged.

Acknowledgment

The University of South Africa is recognized for providing the laboratory space for this study. Biotechnology research group of Prof Atagana is highly acknowledged.

References

- [1] S. Khan, M. Afzal, S. Iqbal, Q. M. Khan, "Plant-bacteria partnerships for the remediation of hydrocarbon contaminated soils," *Chemosphere*, vol. 90, no. 4, pp. 1317–1332, 2014.
- [2] J. Liu, S. Liu, K. Sun, Y. Sheng, Y. Gu, Y. Gao, "Colonization on root surface by a phenanthrene-degrading endophytic bacterium and its application for reducing plant phenanthrene contamination," *Plos/one* vol. 9, no. 9, pp. 1-8, 2014.
- [3] M. Uchino, O. Shida, T. Uchimura, K. Komagata, "Recharacterization of *Pseudomonas fulva* Iizuka and Komagata, and proposals of *Pseudomonas parafulva* sp.nov. and *Pseudomonas cremocricolorata* sp. nov.," *J Gen Appl Microbiol*, vol. 47, no. pp. 247–61, 2001.
- [4] R. O. Anyasi, H. I. Atagana, R. Sutherland, "Identification and characterization of PAH-degrading endophytes isolated from plants growing around a sludge dam," *Int. J. Phytorem*, vol. 21, no. 7, pp. 672-682, 2019.
- [5] B. Van Aken, C. Peres, S. Doty, J. Yoon, J. Schnoor, "*Methylobacterium populi* sp. nov., a novel aerobic, pink-pigmented, facultative methylotrophic, methane-utilizing bacterium isolated from poplar trees (*Populus deltoids x nigra* DN34)," *Evol Microbiol*, vol. 54, no. 4, pp. 1191-1196, 2004.

- [6] N. Weyens, D. van der Lelie, T. Artois, K. Smeets, K. Taghavi, L. Newman, R. Carlier, J Vangronsveld. "Bioaugmentation with engineered endophyte bacteria improves contaminant fate in phytoremediation," *Environ. Sci. Technol.* Vol. 43, no. 24, pp. 9413-9418, 2009a.
- [7] N. Weyens, D. Van Der Lelie, S. Taghavi, L. Newman, J. "Vangronsveld, Exploiting plant-microbe partnerships to improve biomass production and remediation," *Trends Biotechnol.* Vol. 27, no. 10, pp. 591-598, 2009b.
- [8] K. Germaine, X. Liu, G. Cabellos, J. Hogan, D. Ryan, D. N. Dowling, "Bacterial endophyte-enhanced phytoremediation of the organochlorine herbicide 2,4-dichlorophenoxyacetic acid," *FEMS Microbiol Ecol.* 57, no. 2, pp. 302-310, 2010.
- [9] L. E. De Oliveira-Costa, M. V. De Queiroz, A. C. Borges, C. A. De Moraes, E. F. De Araújo, "Isolation and characterization of endophytic bacteria isolated from the leaves of the common bean (*Phaseolus vulgaris*)," *Braz J Microbiol.* vol. 43, no.4, pp. 1562–1575, 2012.
- [10] A. Galazka, R. Galazka, "Phytoremediation of Polycyclic Aromatic Hydrocarbons in Soils Artificially Polluted Using Plant-Associated-Endophytic Bacteria and *Dactylis glomerata* as the Bioremediation Plant," *Pol J Microbiol.* 64, no. 3, pp. 241-253, 2015.
- [11] Anyasi R. O, Atagana H. I. Profiling of plants at petroleum contaminated site for phytoremediation. *Int J Phytorem.* vol. 20, no. 4, pp. 352-364. 2018.
- [12] H. I. Atagana, Bioremediation of co-contamination of crude oil and heavy metals soil by phytoremediation using *Chromolaena odorata* (L) King & H.E. Robinson. *Water Air Soil Poll.* vol.215, no. 1-4, pp. 261-271, 2011b.
- [13] C. L. Rugh. "Mercury detoxification with endophytic plants and other biotechnological breakthroughs for phytoremediation," *In Vitro Cell Dev. Biol. Plant.* vol 37, no. 3, pp. 321-325, 2001.
- [14] S. P. Bizily, C.L. Rugh, R.B. Meagher, Phytodetoxification of hazardous organomercurials by genetically engineered plants. *Nat Biotechnol.* Vol. 18, no. 2 pp. 213-217, 2000.
- [15] R. O. Anyasi, Bioremediation of polychlorinated biphenyls (PCB)-contaminated soil by phytoremediation with *Chromolaena odorata* (L) King and Robinson. MSc thesis, University of South Africa, pp. 50-169, 2012.
- [16] P. Tanhan, M. Kruatrachue, P. Pokethitiyook, R. Chaiyarat, "Uptake and accumulation of cadmium, lead, and zinc by Siam weed [*Chromolaena odorata* (L.) King & Robinson]," *Chemos.* vol. 68, no. 2, pp. 323–329, 2007.
- [17] S. Singh, V. Thorat, C. P. Kaushik, K. Raj, S. Eapen, S. F. D'Souza, "Potential of *Chromolaena odorata* for phytoremediation of ¹³⁷Cs from solution and low level nuclear waste," *J Haz Mat.* vol. 162, no. 2-3, pp. 743-745, 2009.
- [18] H. I. Atagana, "The potential of *Chromolaena odorata* (L) to decontaminate used engine oil impacted soil under greenhouse conditions," *Int J Phytorem.* vol. 13, no. 7, pp. 627-641. 2011a.
- [19] M. I. Russo S. A. Pelizza, M. N. Cabello, S.A. Stenglein, A.C. Scorsetti, "Endophytic colonization of tobacco, corn, wheat and soybeans by the fungal entomopathogen *Beauveria bassiana* (Ascomycota, Hypocreales)," *Biocontrol Sci. Tech.* 25, no. 4, pp. 475-480. 2015.
- [21] D. B. Sauer, R Burroughs, "Disinfection of seed surfaces with Sodium Hypochlorite," *Phytopathology.* vol. 76, no. 3, pp. 745-749, 1986.
- [22] G. Berg, K. Smalla, "Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere," *FEMS Microbiol Ecol.* vol. 68, no. 1, pp. 1–13, 2009.
- [23] K. J. Umar, M. J. Muhammad, N. A. Sani, S. Muhammad, M.T. Umar, "Comparative Study of Antioxidant Activities of the Leaves and Stem of *Ipomoea aquatica* Forsk (Water Spinach)," *Niger J Basic Appl Sci.* 23, no. 1, pp. 81–84, 2017.
- [24] K. J. Germaine, X. Liu, G. G. Cabellos, J .P. Hogan, D. Ryan, D. N. Dowling, "Bacterial endophyte-enhanced phytoremediation of the organochlorine herbicide 2,4-dichlorophenoxyacetic acid," *FEMS Microbiol Ecol.* vol. 57, no. 2, pp. 302–310, 2009.
- [25] E. D. Mitter, R., Kataoka, J. R. de Freitas, J. J. Germida, „Potential use of endophytic root bacteria and host plants to degrade hydrocarbons," *Inter J Phytorem.* vol. 21, no. 2, pp. 1-11, 2019.
- [26] S. Eskandary, A. Tahmourespour, M. Hoodaji, A. Abdollahi, "The synergistic use of plant and isolated bacteria to clean up polycyclic aromatic hydrocarbons from contaminated soil," *J. Environ Health Sci. Eng.* vol. 15 no.12, pp. 1-8, . 2017.
- [27] D. Jain, S. Kachhwaha, R. Jain, S. L. Kothari, "PCR based detection of cry genes in indigenous strains of *Bacillus thuringiensis* isolated from the soils of Rajasthan," *Indian J Biotechnol.* vol. 11, no. 4, pp. 491-494, 2012.